

## A statistical analysis of three ensembles of crop model responses to temperature and CO<sub>2</sub> concentration

D. Makowski<sup>a,\*</sup>, S. Asseng<sup>b</sup>, F. Ewert<sup>c</sup>, S. Bassu<sup>a</sup>, J.L. Durand<sup>d</sup>, T. Li<sup>e</sup>, P. Martre<sup>f,g</sup>, M. Adam<sup>h</sup>, P.K. Aggarwal<sup>i</sup>, C. Angulo<sup>c</sup>, C. Baron<sup>j</sup>, B. Basso<sup>k</sup>, P. Bertuzzi<sup>l</sup>, C. Biernath<sup>m</sup>, H. Boogaard<sup>n</sup>, K.J. Boote<sup>o</sup>, B. Bouman<sup>e</sup>, S. Bregaglio<sup>p</sup>, N. Brisson<sup>a,l,1</sup>, S. Buis<sup>q</sup>, D. Cammarano<sup>pp</sup>, A.J. Challinor<sup>r,s,t</sup>, R. Confalonieri<sup>p</sup>, J.G. Conijn<sup>u</sup>, M. Corbeels<sup>v,w</sup>, D. Deryng<sup>x</sup>, G. De Sanctis<sup>rr</sup>, J. Doltra<sup>y</sup>, T. Fumoto<sup>z</sup>, D. Gaydon<sup>A</sup>, S. Gayler<sup>B</sup>, R. Goldberg<sup>C</sup>, R.F. Grant<sup>D</sup>, P. Grassini<sup>E</sup>, J.L. Hatfield<sup>F</sup>, T. Hasegawa<sup>z</sup>, L. Heng<sup>G</sup>, S. Hoek<sup>n</sup>, J. Hooker<sup>H</sup>, L.A. Hunt<sup>I</sup>, J. Ingwersen<sup>J</sup>, R.C. Izaurralde<sup>K,kk</sup>, R.E.E. Jongschaap<sup>v,w</sup>, J.W. Jones<sup>b</sup>, R.A. Kemanian<sup>L</sup>, K.C. Kersebaum<sup>M</sup>, S.-H. Kim<sup>N</sup>, J. Lizaso<sup>O</sup>, M. Marcaida III<sup>e</sup>, C. Müller<sup>P</sup>, H. Nakagawa<sup>Q</sup>, S. Naresh Kumar<sup>R</sup>, C. Nendel<sup>M</sup>, G.J. O'Leary<sup>S</sup>, J.E. Olesen<sup>T</sup>, P. Oriol<sup>h</sup>, T.M. Osborne<sup>U</sup>, T. Palosuo<sup>ff</sup>, M.V. Pravia<sup>L,V</sup>, E. Priesack<sup>m</sup>, D. Ripoche<sup>I</sup>, C. Rosenzweig<sup>C</sup>, A.C. Ruane<sup>C</sup>, F. Ruget<sup>q,w</sup>, F. Sau<sup>X</sup>, M.A. Semenov<sup>Y</sup>, I. Shcherbak<sup>k</sup>, B. Singh<sup>ss</sup>, U. Singh<sup>Z</sup>, H.K. Soo<sup>aa</sup>, P. Steduto<sup>bb</sup>, C. Stöckle<sup>cc</sup>, P. Stratonovitch<sup>Y</sup>, T. Streck<sup>J</sup>, I. Supit<sup>dd</sup>, L. Tang<sup>ee</sup>, F. Tao<sup>ff</sup>, E.I. Teixeira<sup>gg</sup>, P. Thorburn<sup>hh</sup>, D. Timlin<sup>ii</sup>, M. Travasso<sup>jj</sup>, R.P. Rötter<sup>ff</sup>, K. Waha<sup>P,qq</sup>, D. Wallach<sup>kk</sup>, J.W. White<sup>ll</sup>, P. Wilkens<sup>Z</sup>, J.R. Williams<sup>mm</sup>, J. Wolf<sup>dd</sup>, X. Yin<sup>nn</sup>, H. Yoshida<sup>Q</sup>, Z. Zhang<sup>oo</sup>, Y. Zhu<sup>ee</sup>

<sup>a</sup> INRA, UMR 211 Agronomie INRA AgroParisTech, 78850 Thiverval-Grignon, France

<sup>b</sup> Agricultural & Biological Engineering Department, University of Florida, Gainesville, FL 32611, USA

<sup>c</sup> Institute of Crop Science and Resource Conservation INRES, University of Bonn, 53115, Germany

<sup>d</sup> Unité de Recherche Pluridisciplinaire sur la Prairie et les Plantes Fourragères, INRA, CS 80006, 86600 Lusignan, France

<sup>e</sup> International Rice Research Institute, Los Baños, Philippines

<sup>f</sup> INRA, UMR1095 Genetic, Diversity and Ecophysiology of Cereals (GDEC), F-63 100 Clermont-Ferrand, France

<sup>g</sup> Blaise Pascal University, UMR1095 GDEC, F-63 170 Aubière, France

<sup>h</sup> CIRAD, UMR AGAP/PAM, Av. Agropolis, Montpellier, France

<sup>i</sup> CGIAR Research program on Climate Change, Agriculture and Food Security, International Water Management Institute, New Delhi 110012, India

<sup>j</sup> CIRAD, UMR TETIS, 500 rue J-F. Breton, Montpellier F-34093, USA

<sup>k</sup> Department of Geological Sciences and W.K. Kellogg Biological Station, Michigan State University East Lansing, MI 48823, USA

<sup>l</sup> INRA, US1116 AgroClim, F-84 914 Avignon, France

<sup>m</sup> Institute of Soil Ecology, Helmholtz Zentrum München – German Research Center for Environmental Health, Neuherberg D-85764, Germany

<sup>n</sup> Centre for Geo-Information, Alterra, P.O. Box 47, 6700AA Wageningen, The Netherlands

<sup>o</sup> Department of Agronomy, University of Florida, Gainesville, FL 32611-0500, USA

<sup>p</sup> Cassandra lab, University of Milan, Milan, Italy

\* Corresponding author.

E-mail addresses: [makowski@grignon.inra.fr](mailto:makowski@grignon.inra.fr) (D. Makowski), [sasseng@ufl.edu](mailto:sasseng@ufl.edu) (S. Asseng), [fewert@uni-bonn.de](mailto:fewert@uni-bonn.de) (F. Ewert), [jean-louis.durand@lusignan.inra.fr](mailto:jean-louis.durand@lusignan.inra.fr) (J.L. Durand), [t.li@irri.org](mailto:t.li@irri.org) (T. Li), [pierre.martre@clermont.inra.fr](mailto:pierre.martre@clermont.inra.fr) (P. Martre), [p.k.aggarwal@cigar.org](mailto:p.k.aggarwal@cigar.org) (P.K. Aggarwal), [klav@uni-bonn.de](mailto:klav@uni-bonn.de) (C. Angulo), [basso@msu.edu](mailto:basso@msu.edu) (B. Basso), [dominique.ripoche@avignon.inra.fr](mailto:dominique.ripoche@avignon.inra.fr) (P. Bertuzzi), [priesack@helmholtz-muenchen.de](mailto:priesack@helmholtz-muenchen.de) (C. Biernath), [kjboote@ufl.edu](mailto:kjboote@ufl.edu), [pkaggarwal.iari@gmail.com](mailto:pkaggarwal.iari@gmail.com) (K.J. Boote), [simone.bregaglio@unimi.it](mailto:simone.bregaglio@unimi.it) (S. Bregaglio), [patrick.bertuzzi@avignon.inra.fr](mailto:patrick.bertuzzi@avignon.inra.fr) (N. Brisson), [davide.cammarano@hutton.ac.uk](mailto:davide.cammarano@hutton.ac.uk) (D. Cammarano), [aj.challinor@leeds.ac.uk](mailto:aj.challinor@leeds.ac.uk) (A.J. Challinor), [roberto.confalonieri@unimi.it](mailto:roberto.confalonieri@unimi.it) (R. Confalonieri), [sjaak.conijn@wur.nl](mailto:sjaak.conijn@wur.nl), [raymond.jongschaap@wur.nl](mailto:raymond.jongschaap@wur.nl) (J.G. Conijn), [corbeels@cirad.fr](mailto:corbeels@cirad.fr) (M. Corbeels), [giacomode-santis@jrc.ec.europa.eu](mailto:giacomode-santis@jrc.ec.europa.eu) (G. De Sanctis), [jordidoltra@cifacantabria.org](mailto:jordidoltra@cifacantabria.org) (J. Doltra), [Don.Gaydon@csiro.au](mailto:Don.Gaydon@csiro.au) (D. Gaydon), [Sebastian.gayler@uni-tuebingen.de](mailto:Sebastian.gayler@uni-tuebingen.de) (S. Gayler), [cynthia.rosenzweig@nasa.gov](mailto:cynthia.rosenzweig@nasa.gov) (R. Goldberg), [rgrant@ualberta.ca](mailto:rgrant@ualberta.ca) (R.F. Grant), [jerry.hatfield@ars.usda.gov](mailto:jerry.hatfield@ars.usda.gov) (J.L. Hatfield), [L.Heng@iaea.org](mailto:L.Heng@iaea.org) (L. Heng), [j.hooker@reading.ac.uk](mailto:j.hooker@reading.ac.uk) (J. Hooker), [thunt@uoguelph.ca](mailto:thunt@uoguelph.ca) (L.A. Hunt), [joachim.ingwersen@uni-hohenheim.de](mailto:joachim.ingwersen@uni-hohenheim.de) (J. Ingwersen), [cizaurra@umd.edu](mailto:cizaurra@umd.edu) (R.C. Izaurralde), [sjaak.conijn@wur.nl](mailto:sjaak.conijn@wur.nl) (R.E.E. Jongschaap), [jimj@ufl.edu](mailto:jimj@ufl.edu) (J.W. Jones), [vpravia@inia.org.uv](mailto:vpravia@inia.org.uv) (R.A. Kemanian), [ckersebaum@zalf.de](mailto:ckersebaum@zalf.de) (K.C. Kersebaum), [christoph.mueller@pik-potsdam.de](mailto:christoph.mueller@pik-potsdam.de) (C. Müller), [nareshkumar.soora@gmail.com](mailto:nareshkumar.soora@gmail.com) (S. Naresh Kumar), [nendel@zalf.de](mailto:nendel@zalf.de) (C. Nendel), [garry.O'leary@ecodev.vic.gov.au](mailto:garry.O'leary@ecodev.vic.gov.au) (G.J. O'Leary), [jeo@agro.au.dk](mailto:jeo@agro.au.dk) (J.E. Olesen), [t.m.osborne@reading.ac.uk](mailto:t.m.osborne@reading.ac.uk) (T.M. Osborne), [taru.palosuo@luke.fi](mailto:taru.palosuo@luke.fi) (T. Palosuo), [christian.biernath@helmholtz-muenchen.de](mailto:christian.biernath@helmholtz-muenchen.de) (E. Priesack), [ccrag1@yahoo.com](mailto:ccrag1@yahoo.com) (C. Rosenzweig), [alexander.c.ruane@nasa.gov](mailto:alexander.c.ruane@nasa.gov) (A.C. Ruane), [mikhail.semenov@rothamsted.ac.uk](mailto:mikhail.semenov@rothamsted.ac.uk) (M.A. Semenov), [shcherba@mail.msu.edu](mailto:shcherba@mail.msu.edu) (I. Shcherbak), [Balwinder.singh@cgiar.org](mailto:Balwinder.singh@cgiar.org) (B. Singh), [Pasquale.Steduto@fao.org](mailto:Pasquale.Steduto@fao.org) (P. Steduto), [stockle@wsu.edu](mailto:stockle@wsu.edu) (C. Stöckle), [pierre.stratonovitch@rothamsted.ac.uk](mailto:pierre.stratonovitch@rothamsted.ac.uk) (P. Stratonovitch), [tstreck@uni-hohenheim.de](mailto:tstreck@uni-hohenheim.de) (T. Streck), [iwan.supit@wur.nl](mailto:iwan.supit@wur.nl) (I. Supit), [tangl@njau.edu.cn](mailto:tangl@njau.edu.cn) (L. Tang), [fulu.tao@luke.fi](mailto:fulu.tao@luke.fi) (F. Tao), [Edmar.teixeira@plantandfood.co.nz](mailto:Edmar.teixeira@plantandfood.co.nz) (E.I. Teixeira), [peter.thorburn@csiro.au](mailto:peter.thorburn@csiro.au) (P. Thorburn), [mtravasso@cnia.inta.gov.ar](mailto:mtravasso@cnia.inta.gov.ar) (M. Travasso), [reimund.rotter@luke.fi](mailto:reimund.rotter@luke.fi) (R.P. Rötter), [katharina.waha@csiro.au](mailto:katharina.waha@csiro.au) (K. Waha), [daniel.wallach@toulouse.inra.fr](mailto:daniel.wallach@toulouse.inra.fr) (D. Wallach), [jeffrey.white@ars.usda.gov](mailto:jeffrey.white@ars.usda.gov) (J.W. White), [jwilliams@brc.tamus.edu](mailto:jwilliams@brc.tamus.edu) (J.R. Williams), [joost.wolf@wur.nl](mailto:joost.wolf@wur.nl) (J. Wolf), [yanzhu@njau.edu.cn](mailto:yanzhu@njau.edu.cn) (Y. Zhu).

<sup>1</sup> # Dr. Nadine Brisson passed away in 2011 while this work was being carried out.

- <sup>q</sup> INRA, UMR1114 EMMAH, F-84914 Avignon, France
- <sup>r</sup> Institute for Climate and Atmospheric Science, School of Earth and Environment, University of Leeds, Leeds LS29JT, UK
- <sup>s</sup> CGIAR-ESSP Program on Climate Change, Agriculture and Food Security, International Centre for Tropical Agriculture (CIAT), A.A. 6713 Cali, Colombia
- <sup>t</sup> CGIAR-ESSP Program on Climate Change, Agriculture and Food Security, International Centre for Tropical Agriculture (CIAT), A.A. 6713 Cali, Colombia
- <sup>u</sup> WUR-Plant Research International, Wageningen University and Research Centre, Droeveendaalsesteeg 1, 6708PB Wageningen, The Netherlands
- <sup>v</sup> Agro-ecology and Sustainable Intensification of Annual Crops, CIRAD, Avenue Agropolis, 34398 Montpellier Cedex 5, France
- <sup>w</sup> Embrapa-Cerrados, PO Box 8233, 73301-970 Planaltina, DF, Brazil
- <sup>x</sup> Tyndall Centre for Climate Change research and School of Environmental Sciences, University of East Anglia, Norwich NR4 7TJ, UK
- <sup>y</sup> Cantabrian Agricultural Research and Training Centre (CITA), Cantabria Government, 39600 Muriedas, Spain
- <sup>z</sup> National Institute for Agro-Environmental Sciences, Tsukuba, Ibaraki, 305-8604, Japan
- <sup>A</sup> CSIRO Agriculture Flagship, Dutton Park, QLD 4102, Australia
- <sup>B</sup> WESS-Water & Earth System Science Competence Cluster, University of Tübingen, 72074 Tübingen, Germany
- <sup>C</sup> NASA Goddard Institute for Space Studies, New York, NY 10025, USA
- <sup>D</sup> Department of Renewable Resources, University of Alberta, Edmonton, AB, Canada T6G 2E3
- <sup>E</sup> Department of Agronomy and Horticulture, University of Nebraska-Lincoln, 178 Keim Hall-East Campus, Lincoln, NE 68503-0915, USA
- <sup>F</sup> National Laboratory for Agriculture and Environment, Ames, IA 50011, USA
- <sup>G</sup> IAEA, Vienna, Austria
- <sup>H</sup> Agriculture Department, University of Reading, Reading RG66AR, UK
- <sup>I</sup> Department of Plant Agriculture, University of Guelph, Guelph, Ontario, Canada N1G 2W1
- <sup>J</sup> Institute of Soil Science and Land Evaluation, Universität Hohenheim, 70599 Stuttgart, USA
- <sup>K</sup> Department of Geographical Sciences, University of Maryland, College Park, MD 20742, USA
- <sup>L</sup> Instituto Nacional de Investigación Agropecuaria (INIA), Address: Ruta 8 km 281, Treinta y Tres, Uruguay 33000, USA
- <sup>M</sup> Institute of Landscape Systems Analysis, Leibniz Centre for Agricultural Landscape Research, 15374 Müncheberg, Germany
- <sup>N</sup> School of Environmental and Forest Sciences, College of the Environment, University of Washington, Seattle, WA, USA
- <sup>O</sup> Department Producción Vegetal, Fitotecnia, Univ. Politécnica de Madrid, 28040 Madrid, Spain
- <sup>P</sup> Potsdam Institute for Climate Impact Research, 14473 Potsdam, Germany
- <sup>Q</sup> National Agriculture and Food Research Organization, Japan
- <sup>R</sup> Centre for Environment Science and Climate Resilient Agriculture, Indian Agricultural Research Institute, IARI PUSA, New Delhi 110 012, India
- <sup>S</sup> Grains Innovation Park, Department of Economic Development Jobs, Transport and Resources, Horsham 3400, Australia
- <sup>T</sup> Department of Agroecology, Aarhus University, 8830 Tjele, Denmark
- <sup>U</sup> NCAS-Climate, Walker Institute, University of Reading, RG6 6BB, UK
- <sup>V</sup> Department of Plant Science, The Pennsylvania State University, 247 Agricultural Sciences and Industries Building, University Park, PA 16802, USA
- <sup>W</sup> UAPV, UMR1114 EMMAH, F-84914 Avignon, France
- <sup>X</sup> Department Biología Vegetal, Univ. Politécnica de Madrid, 28040 Madrid, Spain
- <sup>Y</sup> Computational and Systems Biology Department, Rothamsted Research, Harpenden, Herts AL5 2JQ, UK
- <sup>Z</sup> International Fertilizer Development Institute, Florence, AL, USA
- <sup>aa</sup> School of Environmental and Forest Sciences, College of the Environment, University of Washington, Seattle, WA, USA
- <sup>bb</sup> FAO, Rome, Italy
- <sup>cc</sup> Biological Systems Engineering, Washington State University, Pullman, WA 99164-6120, USA
- <sup>dd</sup> Plant Production Systems & Earth System Science, Wageningen University, 6700AA Wageningen, The Netherlands
- <sup>ee</sup> National Engineering and Technology Center for Information Agriculture, Nanjing Agricultural University, China
- <sup>ff</sup> Natural Resources Institute Finland (Luke), FI-00790 Helsinki, Finland
- <sup>gg</sup> Sustainable Production, The New Zealand Institute for Plant & Food Research Limited, Lincoln, Canterbury, New Zealand
- <sup>hh</sup> CSIRO Agriculture Flagship, Dutton Park, QLD 4102, Australia
- <sup>ii</sup> USDA/ARS, Crop Systems and Global Change Laboratory, 10300 Baltimore avenue, BLDG 001 BARC-WEST, Beltsville, MD 20705-2350, USA
- <sup>jj</sup> Institute for Climate and Water, INTA-CIRN, 1712 Castelar, Argentina
- <sup>kk</sup> INRA, UMR 1248 Agrosystèmes et développement territorial (AGIR), 31326 Castanet-Tolosan Cedex, France
- <sup>ll</sup> Arid-Land Agricultural Research Center, Maricopa, AZ 85138, USA
- <sup>mmm</sup> Texas AgriLife Research and Extension, Texas A&M University, USA
- <sup>nn</sup> Centre for Crop Systems Analysis, Wageningen University, The Netherlands
- <sup>oo</sup> State Key Laboratory of Earth Surface Processes and Resource Ecology, Beijing Normal University, Beijing 100875, China
- <sup>pp</sup> The James Hutton Institute Invergowrie, Dundee DD2 5DA, Scotland, UK
- <sup>qq</sup> CSIRO Agriculture, 306 Carmody Road, 4067 St. Lucia, QLD, Australia
- <sup>rr</sup> European Commission Joint Research Center, via Enrico Fermi, 2749, Ispra 21027, Italy
- <sup>ss</sup> CIMMYT-India, Aggarwal Corporate Towers, New Delhi 110 008, India

## ARTICLE INFO

## Article history:

Received 6 March 2015

Received in revised form 29 July 2015

Accepted 20 September 2015

## Keywords:

Climate change

Crop model

Emulator

Meta-model

Statistical model

Yield

## ABSTRACT

Ensembles of process-based crop models are increasingly used to simulate crop growth for scenarios of temperature and/or precipitation changes corresponding to different projections of atmospheric CO<sub>2</sub> concentrations. This approach generates large datasets with thousands of simulated crop yield data. Such datasets potentially provide new information but it is difficult to summarize them in a useful way due to their structural complexities. An associated issue is that it is not straightforward to compare crops and to interpolate the results to alternative climate scenarios not initially included in the simulation protocols. Here we demonstrate that statistical models based on random-coefficient regressions are able to emulate ensembles of process-based crop models. An important advantage of the proposed statistical models is that they can interpolate between temperature levels and between CO<sub>2</sub> concentration levels, and can thus be used to calculate temperature and [CO<sub>2</sub>] thresholds leading to yield loss or yield gain, without re-running the original complex crop models. Our approach is illustrated with three yield datasets simulated by 19 maize models, 26 wheat models, and 13 rice models. Several statistical models are fitted to these datasets, and are then used to analyze the variability of the yield response to [CO<sub>2</sub>] and temperature. Based on our results, we show that, for wheat, a [CO<sub>2</sub>] increase is likely to outweigh the negative effect of a temperature increase of +2 °C in the considered sites. Compared to wheat, required levels of [CO<sub>2</sub>] increase are much higher for maize, and intermediate for rice. For all crops, uncertainties in simulating climate change impacts increase more with temperature than with elevated [CO<sub>2</sub>].

© 2015 Elsevier B.V. All rights reserved.

## 1. Introduction

Many studies have been carried out in recent decades to assess the effects of climate change on crop yield and other key crop characteristics. In these studies, one or several crop models were used to simulate crop growth and development for different projections of atmospheric CO<sub>2</sub> concentration, temperature and precipitation changes (Semenov et al., 1996; Tubiello and Ewert, 2002; White et al., 2011). AgMIP, the Agricultural Model Intercomparison and Improvement Project (Rosenzweig et al., 2013), builds on these studies to explore the value of an ensemble of crop models for assessing effects of climate change scenarios for several crops in contrasting environments.

The AgMIP studies generate large datasets, including thousands of simulated crop yield data. They include series of yield values that are obtained by using standardized protocols that combine several crop models with different climate scenarios defined by several climatic variables (temperature, CO<sub>2</sub>, precipitation, etc.). Such datasets potentially provide new information on the possible effects of different climate change scenarios on crop yields. However, it is difficult to summarize them in a useful way due to their structural complexity; simulated yield data can differ among contrasting climate scenarios, sites, and crop models. Another issue is that it is not straightforward to interpolate the results obtained for the considered scenarios to alternative climate scenarios not considered in the initial simulation protocols. Additional crop model simulations for new climate scenarios is an option but this approach is costly, especially when a large number of crop models is used to generate the simulated data.

Statistical models have been used to analyze responses of measured yield data to climate variables in past studies (Lobell et al., 2011). They were also recently used in meta-analyses on the effect of climate change on crop yields (Wilcox and Makowski, 2014; Challinor et al., 2014). However, the use of a statistical model to analyze the variability of crop model responses to climate change factors is a rather new idea. We demonstrate herewith that statistical methods can play an important role in analyzing simulated yield datasets obtained with ensembles of process-based crop models using standardized protocols. Formal statistical analysis is helpful to estimate the effects of different climatic variables on yield, and to describe the between-model variability of these effects. Statistical methods can also be used to develop meta-models, i.e., statistical models summarizing process-based crop models. Such meta-models may enable scientists to explore more efficiently the effects of new climate change scenarios not initially included in the simulation protocol.

Our approach is illustrated with three datasets of simulated yields obtained by AgMIP for maize, wheat, and rice generated by ensembles of process-based crop models (Asseng et al., 2013; Bassu et al., 2014; Li et al., 2015). The yield datasets were used to develop a meta-model that provides a simplified representation of the original ensembles of crop models. The proposed meta-model is a statistical regression with random coefficients describing the variability of the simulated yield data across the original crop models. Once fitted to the simulated yield datasets, the meta-models were used to analyze the variability of the projected effects of climate changes among crop models, and between alternative crops. The meta-models were also used to study the effects of temperature-change and CO<sub>2</sub>-change scenarios that were not initially tested with the original ensemble of crop models. Finally, the results obtained with the meta-model were used to compare simulated uncertainties and to assess the impact of temperature and CO<sub>2</sub> concentration changes on yields of maize, wheat, and rice.

## 2. Materials and methods

### 2.1. Simulated yield data

We used the maize, wheat, and rice datasets presented by Asseng et al. (2013), Bassu et al. (2014), and Li et al. (2015). Yield data were simulated with 19 maize models, 26 wheat models, and 13 rice models. For each crop species, models were calibrated and then run for four contrasting sites located in France (Lusignan), USA (Ames), Brazil (Rio Verde), and Tanzania (Morogoro) for maize, in The Netherlands (Wageningen), Argentina (Balcarce), India (New Delhi), and Australia (Wongan Hills) for wheat, and in the Philippines (Los Baños), China (Nanjing), India (Ludhiana) and Japan (Shizukuishi) for rice.

The simulation protocols and climate scenarios are described in Rosenzweig et al. (2013), Asseng et al. (2013), Bassu et al. (2014), and Li et al. (2015). The baseline scenario corresponded to the 1980–2010 historical climates and assumed a CO<sub>2</sub> concentration of 360 ppm (mean of 1995). The other climate scenarios were defined from the baseline weather series by changing the daily maximum and minimum temperature and CO<sub>2</sub> concentration. For all species, four temperature changes (+0, +3, +6, +9°C) and five atmospheric CO<sub>2</sub> concentration changes (+0, +90, +180, +270, +360 ppm) were used. Thirty years of yield data were generated with each crop model for each scenario, and the simulated yield values were averaged over the years. The total number of mean yield data was equal to 1764 (441 per site) for maize, to 2592 (648 per site) for wheat, to 1138 (282–286 per site) for rice. Details of the maize, wheat, and rice protocols can be found in Bassu et al. (2014), Asseng et al. (2013), and Li et al. (2015) respectively.

### 2.2. Statistical model

Simulated maize, wheat and rice yield data were analyzed using two-level statistical random-effect models (Davidian and Giltinan, 1995; Pinheiro and Bates, 2000) relating mean yield (averaged over 30 years) to temperature change, atmospheric CO<sub>2</sub> concentration change, and their interaction. The following statistical model was used to analyze yield data for each crop and each site separately:

Level 1, within crop model

$$Y_{ij} = \alpha_{0i} + \alpha_{1i}\Delta T_{ij} + \alpha_{2i}\Delta T_{ij}^2 + \alpha_{3i}\Delta C_{ij} + \alpha_{4i}\Delta C_{ij}^2 + \alpha_{5i}\Delta C_{ij}\Delta T_{ij} + \varepsilon_{ij}, \quad (1)$$

where  $\varepsilon_{ij} \sim N(0, \tau^2)$  (assumed independently and identically distributed),  $Y_{ij}$  is the mean yield (averaged over 30 years) simulated with the  $i$ th crop model,  $i = 1, \dots, P$ , for the  $j$ th scenario;  $j = 1, \dots, Q_i$ ,  $\Delta T_{ij}$ ,  $\Delta C_{ij}$  are the temperature change (compared to the baseline scenario), and atmospheric CO<sub>2</sub> concentration change for model  $i$  and scenario  $j$ ,  $Q_i$  is the number of scenarios tested with model  $i$ ,  $\tau^2$  is a variance describing the residual error,

Level 2, between crop models

$$\alpha_{ki} \sim N(\mu_k, \sigma_k^2), \quad k = 0, \dots, 5. \quad (2)$$

where  $\alpha_{ki}$ ,  $k = 0, \dots, 5$ , are six random regression coefficients distributed according to independent Gaussian probability distributions,  $\mu_k$ ,  $k = 0, \dots, 5$ , are the seven mean regression coefficient values representing the mean yield baseline ( $\mu_0$ ), the mean effect of temperature, the mean effect of CO<sub>2</sub>, and the mean effect of temperature-CO<sub>2</sub> interaction ( $\mu_1 \dots \mu_5$ ) over the  $P$  crop models (i.e., the expected values of  $\alpha_{ki}$ ,  $k = 0, \dots, 5$ ). The six variances,  $\sigma_k^2$ ,  $k = 0, \dots, 5$ , describe the between-model variability of the random regression coefficients (i.e., the variances of  $\alpha_{ki}$ ,  $k = 0, \dots, 5$ ).

This statistical model assumes that the ensemble of  $P$  crop models is a sample taken within a population including all possible crop models for a given crop while flexibly allowing for the incorporation of additional crop models in the future. The probability distributions defined by Eq. (2) describe the between-crop model variability of the yield response to climate change factors within the whole population of crop models. These probability distributions cover the ranges of climate effects considered by different crop models. The relationship defined by Eq. (1) is assumed to be valid for all crop models, but its parameters  $\alpha_{ki}$ ,  $k=0, \dots, 5$ , are assumed to vary among crop models. This statistical model describes 30-year mean yield responses and is not intended to describe the year-to-year variability of crop yields. Considering year-to-year variability would require extra random terms and additional parameters and would overly complicate the calculated model. This option was thus not considered here. The statistical model could be easily extended to deal with additional variables such as rainfall or farmers' practices.

The population parameters of the statistical model  $\mu_k$ ,  $\sigma_k^2$ , and  $\tau^2$  were estimated by restricted maximum likelihood. The model-specific regression coefficients  $\alpha_{ki}$ ,  $k=0, \dots, 5$ ,  $i=1, \dots, P$ , were estimated by Best Linear Unbiased Predictor using the R software package "nlme" (Pinheiro and Bates, 2000), and the estimated values will be henceforth referred to as  $\alpha_{eki}$ . The model was fitted to data for each crop and each site separately, but for all crop models together. Results were analyzed site by site.

### 2.3. Assessment of the statistical model

The statistical model (Eqs. (1) and (2)) was compared to other statistical models, including models with fewer explanatory variables, models with fewer random coefficients, and a model including no random coefficient (i.e., classical linear regression). All models were compared by using the Akaike Information Criterion (AIC, Akaike, 1973), where a lower AIC value corresponds with a better model. The AIC was calculated using the "AIC" function of R. We found that the model defined by Eqs. (1) and (2) led to lower AICs than the simpler models. The AIC of classical linear regression model was very different from the value obtained with the random coefficient model (Eqs. (1) and (2)); the values of AIC obtained with the classical linear regression model were higher by 78–357% depending on the crop and on the site. The assumption that the residual errors  $\varepsilon_{ij}$  were independent was assessed by developing another statistical model that incorporated the correlated residual errors. This model was fitted using the "correlation" argument of the "lme" function of R. The AIC of this model was higher, and the estimated correlation coefficients were very close to zero (from  $-9.9 \times 10^{-3}$  to  $7.6 \times 10^{-4}$ , depending on the crop and on the site). In order to check the assumption of constant residual variance, a statistical model with a non-constant residual variance was fitted to the data using the "weights" argument of the "lme" function of R. This model was not selected because its AIC was higher. The quality of fit of the statistical model (Eqs. (1) and (2)) was also assessed using graphical analysis and by calculating the coefficient of determination ( $R^2$ ). The value of  $R^2$  was 0.99 for all crops (Fig. 1). Outputs of the statistical model (Eqs. (1) and (2)) were also compared to the original simulated yield data in Fig. 2 for three sites and three crop models per site.

### 2.4. Estimation of the effect of climate change on yield

The statistical model described above was used to compute three different types of outputs:




Fig. 1. Simulated values of maize (A), wheat (B), and rice (C) yields as a function of fitted values obtained with the selected statistical models. Each graphic includes yield data simulated in four sites.


- The average yield loss/gain due to climate change over the ensemble of crop models.
- The yield gain/loss estimated for individual crop models due to changes in climate variables.
- The probability of yield loss compared to the baseline yield.

For maize, the average yield difference obtained between a given climate change scenario (characterized by  $\Delta T$  and  $\Delta C$ ) and the baseline scenario was expressed as

$$\Delta Y = \mu_1 \Delta T + \mu_2 \Delta T^2 + \mu_3 \Delta C + \mu_4 \Delta C^2 + \mu_5 \Delta C \Delta T \quad (3)$$

The yield difference described in Eq. (3) is averaged over all crop models; this difference corresponds to an average yield gain or to an average yield loss over the  $P$  crop models. Eq. (3) defines a meta-model that simulates the average output of the original ensemble of crop models. This meta-model enables the computation of the yield differences for any change in temperature and  $\text{CO}_2$ ,  $\Delta T$  and  $\Delta C$ , at each of the four considered sites.





**Fig. 2.** Examples of yield data simulated by three different crop models for maize in Morogoro (Tanzania) (A), for wheat in Wageningen (The Netherlands) (B), and for rice in Shizukuishi (Japan) (C) for three different CO<sub>2</sub> concentrations and a temperature increase of +3 °C. Points of different colors indicate yield data simulated by using different crop models. Curves correspond to the yield values obtained with the statistical models.

For a given crop model  $i$ , the expected yield difference was expressed as

$$\Delta Y_i = \alpha_{e1i} \Delta T + \alpha_{e2i} \Delta T^2 + \alpha_{e3i} \Delta C + \alpha_{e4i} \Delta C^2 + \alpha_{e5i} \Delta C \Delta T \quad (4)$$

Eq. (4) defines a meta-model simulating the output of the  $i$ th crop model. The yield difference (4) is crop model-specific, and represents an estimation of the expected yield difference resulting from changes in temperature and CO<sub>2</sub> concentration change equal to  $\Delta T$  and  $\Delta C$ , calculated with the  $i$ th crop model. It corresponds to an emulation of the mean climate change effect on yield that would have been obtained with the  $i$ th crop model if this crop model was run for a climate change scenario characterized by  $\Delta T$  and  $\Delta C$ .

The statistical model defined by Eqs. (1) and (2) was also used to compute the probability of yield loss  $\text{Prob}(\Delta Y_i > 0)$  that results from a change in the temperature and CO<sub>2</sub> concentration. This

probability was computed from the following Gaussian probability distribution  $N(\mu_{\Delta Y}, \sigma_{\Delta Y}^2)$  as

$$\mu_{\Delta Y} = \mu_1 \Delta T + \mu_2 \Delta T^2 + \mu_3 \Delta C + \mu_4 \Delta C^2 + \mu_5 \Delta C \Delta T \quad (5)$$

$$\sigma_{\Delta Y}^2 = \sigma_1^2 \Delta T^2 + \sigma_2^2 \Delta T^4 + \sigma_3^2 \Delta C^2 + \sigma_4^2 \Delta C^4 + \sigma_5^2 \Delta C^2 \Delta T^2 \quad (6)$$

Note that the variance defined by Eq. (6) is not constant but varies as a function of the climate-scenario characteristics. For illustration, the quantities defined by Eqs. (3)–(6) were calculated for values of  $\Delta T$  and of  $\Delta C$  ranging from 0 to +6 °C (with a step of 0.1 °C) and from 0 to +360 ppm (with a step of 1 ppm) respectively, i.e.,  $\Delta T = 0, 0.1, 0.2, \dots, 5.9, 6$  °C,  $\Delta C = 0, 1, \dots, 359, 360$  ppm. Some of the considered values of  $\Delta T$  and of  $\Delta C$  were initially included in the simulation protocol (e.g.,  $\Delta T = +3$  °C,  $\Delta C = +180$  ppm) but most of them were not (e.g.,  $\Delta T = +2$  °C,  $\Delta T = +4$  °C,  $\Delta C = +100$  ppm). These calculations were done to demonstrate the capability of the meta-model to study the effects of temperature-change and CO<sub>2</sub>-change scenarios that were not initially tested with the original ensemble of crop models.


### 3. Results

#### 3.1. Yield response to increase in temperature

Fig. 3 shows the change in yield from the baseline for one maize site (Fig. 3A), one wheat site (Fig. 3C), and one rice site (Fig. 3E) as affected by an atmospheric CO<sub>2</sub> concentration increase of 180 ppm ([CO<sub>2</sub>] = 540 ppm) and an increase of mean seasonal temperature ranging from 0 °C to 6 °C. Each emulated model yield response is calculated by using the crop model-specific coefficients  $\alpha_{eki}$  ( $k = 0, \dots, 5$ ,  $i = 1, \dots, P$ ) and is plotted with a gray line, and thus can be seen as a substitute for a given crop model, but without the need for re-running the original, process-based crop models. Positive yield differences can be interpreted as mean yield gain and, conversely, negative yield differences can be interpreted as mean yield loss. The solid red curve indicates the mean of the emulated yield responses to the given climate scenario as compared to the baseline, i.e., the effect averaged over all crop models. The red dashed curves indicate the 10th and 90th percentiles of the climate-change effect. About 10% of the crop models predict yield effects lower/higher than the values given by the lower/upper dashed curve.

According to Fig. 3(A, C and E), most crop models estimate that a temperature increase negatively impacts yields of maize, wheat, and rice at these sites. But this effect is highly variable among crop models, with some models predicting little response to temperature. For maize, Fig. 3A illustrates how, on average across the ensemble of crop models, the statistical model emulates a yield loss when the temperature exceeds +1 °C with a CO<sub>2</sub> concentration increase of 180 ppm in Morogoro, Tanzania. In contrast, the models suggest that wheat in Wageningen (The Netherlands) and rice in Shizukuishi (Japan) would require a temperature increase of 3.6 °C and 5 °C, respectively, before experiencing a yield loss. For a CO<sub>2</sub> concentration increase of 180 ppm, the averaged emulated projections reported in Fig. 3(A, C and E) indicate that moderate temperature increases could lead to gains in wheat and rice productions in these two locations. However, some of the considered crop models predict a stronger negative impact of temperature for wheat and rice. The 10th percentile of the emulated wheat and rice yield response to temperature is indeed negative when the temperature increase exceeds 1.5 °C (Fig. 3A, C and E).

Fig. 3 also reveals the large variability among crop models and displays how this variability increases as a function of temperature. The differences between the 10th and 90th percentiles are much larger for higher temperature increases, at a given CO<sub>2</sub> concentration. For example, for wheat in Wageningen (The Netherlands), the difference between the 10th and 90th percentiles is lower




**Fig. 3.** Yield responses to temperature change ( $\Delta T$ ) and  $\text{CO}_2$  change ( $\Delta \text{CO}_2$ ) estimated using the statistical models for maize in Morogoro (Tanzania), wheat in Wageningen (The Netherlands), and rice in Shizuishi (Japan). Yield differences are expressed relatively to a baseline defined over 1981–2010. Each gray curve corresponds to a given crop model, and is estimated with a statistical model using the crop model-specific coefficients  $\alpha_{eki}$ ,  $k = 0, \dots, 5$ ,  $i = 1, \dots, P$ . The solid red curve indicates the mean effect on yield of climate change compared to the baseline scenario, i.e., the effect averaged over all crop models. The red dashed curves indicate the 10th and 90th percentiles of climate change effect computed over all crop models. The blue line indicates zero yield difference. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of the article.)

than  $2 \text{ t ha}^{-1}$  when the temperature increase is equal to  $+1^\circ\text{C}$ , but becomes higher than  $4 \text{ t ha}^{-1}$  when the temperature increase reaches  $+4^\circ\text{C}$ . This result indicates that the differences among crop models and therefore the model uncertainties are much larger for high than for small temperature increases.

This result is confirmed by the probability densities  $N(\mu_{\Delta Y}, \sigma_{\Delta Y}^2)$  (Eqs. (3) and (4)) shown in Fig. 4 for each crop (maize, wheat, rice) and all sites. The distributions presented in Fig. 4 describe the variability of simulated yield loss (or yield gain) values among crop models, for a temperature increase of either

$+2$  or  $+4^\circ\text{C}$  at a concurrent  $\text{CO}_2$  concentration increase equal to  $+180 \text{ ppm}$ . In all sites, the distributions are more peaked for a temperature increase of  $+2^\circ\text{C}$  and are flatter for a temperature increase of  $+4^\circ\text{C}$ . This result reveals that the variability among crop models is systematically greater for a large than for a low temperature increase. The difference is particularly important for rice (Fig. 4E and F). These plots also show which regions are most affected by a  $+2$  or  $+4^\circ\text{C}$  temperature increase, especially for rice, showing the already warm Philippines and India as the most affected sites.




**Fig. 4.** Probability distributions of yield loss (or yield gain) resulting from an increase of temperature of  $+2^\circ\text{C}$  (A, C, E) or  $+4^\circ\text{C}$  (B, D, F) for maize, wheat, and rice (four sites per crop). The  $\text{CO}_2$  concentration increase is set to  $+180$  ppm in all cases ( $[\text{CO}_2] = 570$  ppm). Each distribution describes the variability of the yield loss (or gain) values simulated by different crop models.

### 3.2. Yield response to increase in $\text{CO}_2$ concentration

Fig. 3 shows the effect of climate change at one site for maize (Fig. 3B), wheat (Fig. 3D), and rice (Fig. 3F) yields under increasing levels of  $\text{CO}_2$  concentration, ranging from 0 to  $+360$  ppm from the simulated baseline concentration (i.e.,  $360\text{--}720$  ppm) and for a constant temperature increase of  $+2^\circ\text{C}$ . Fig. 3B illustrates for maize how a majority of the crop models for maize in Tanzania predict a yield loss for a temperature increase of temperature increase of  $+2^\circ\text{C}$  and the full range of considered  $\text{CO}_2$  concentration. For maize in this site, the mean curve suggests that the benefits of an increased  $\text{CO}_2$  concentration are small, and do not outweigh the negative effect resulting from an increase of  $+2^\circ\text{C}$ .

The fitted response curves obtained for wheat in Wageningen (The Netherlands) and for rice in Shizukuishi (Japan) (Fig. 3D and F) show a different pattern. Compared to maize in Tanzania, the effect of a  $\text{CO}_2$  concentration increase is stronger for wheat and rice. This result was expected based on literature for crops with C-3 vs C-4 photosynthesis (Hatfield et al., 2011). The effect of  $\text{CO}_2$  is highly variable among crop models; some models have strongly positive slopes over the range of  $\text{CO}_2$  concentrations, whereas others show slopes close to zero. When averaged over crop models, the negative effect of the  $+2^\circ\text{C}$  temperature increase is overcome by the positive  $\text{CO}_2$  effect (yield gain) as soon as the  $\text{CO}_2$  concentration increase reaches 100 ppm in the two considered sites (Fig. 3D and F). However, the 10th percentiles are negative for all tested  $\text{CO}_2$



**Fig. 5.** Yield loss probability as a function of CO<sub>2</sub> concentration (ranging from +0 to +360 ppm compared to a baseline concentration of 360 ppm) for two levels of temperature increase (+2 or +4 °C). Each curve corresponds to one crop (maize, wheat or rice) and one site. Yield loss probabilities correspond to the proportions of crop models predicting a yield loss. The horizontal dashed line corresponds to a probability of yield loss of 0.5.

concentrations for both wheat and rice, and this result reveals that more than 10% of the crop models predict a yield loss compared to the baseline, even for high CO<sub>2</sub> concentrations (Fig. 3D and F).

Fitted rice yield response to CO<sub>2</sub> concentration in Shizukuishi in Japan (Fig. 3F) is similar to wheat response in Wageningen in The Netherlands (Fig. 3D). On average over crop models, the climate change effect on rice yield is positive for the whole range of tested CO<sub>2</sub> concentrations in Shizukuishi. As for wheat, the 10% percentiles of yield effects are negative for all tested CO<sub>2</sub> concentrations, and this result reveals that more than 10% of the crop models predict a yield loss compared to the baseline, even for high CO<sub>2</sub> concentrations (Fig. 3F). Fig. 3B, D and F shows that between-crop model variability tends to increase with CO<sub>2</sub> concentration, but this effect is much smaller than for the effect of temperature (Fig. 3A, C and E). The level of divergence between crop model predictions does not strongly increase in response to CO<sub>2</sub>.

Fig. 2 also illustrates the yield responses to CO<sub>2</sub> concentration increase, but for a limited number of models and for a temperature increase of +3 °C. This confirms that the yield response to CO<sub>2</sub> concentration is stronger for wheat and rice than for maize.

### 3.3. Probability of yield loss

Fig. 5 gives a more general picture of the results. Here, the three crops (maize, wheat, rice), and two temperature changes (+2 °C and +4 °C) are considered for all the sites. The curves displayed in Fig. 5 show the yield loss probability  $\text{Prob}(\Delta Y < 0)$  attributed to two levels of temperature increase (+2° and +4 °C) as a function of increasing increments of CO<sub>2</sub> concentration, ranging from 0 to +360 ppm. The yield loss probability represents the proportion of crop models that predict a yield loss due to changes in temperature and CO<sub>2</sub>



**Table 1**

Levels of increased CO<sub>2</sub> concentration (ppm) required to obtain a probability of maize and wheat yield gain higher than 0.5 (i.e., 50% chance of yield gain). Concentration increase levels were computed for two values of temperature increase (+2 and +4 °C) and four sites per crop. Baseline scenario corresponds to [CO<sub>2</sub>] = 360 ppm.

Site	Temperature change	
	+2 °C	+4 °C
Maize	[CO <sub>2</sub> ]	[CO <sub>2</sub> ]
Brazil (Rio Verde)	>+360	>+360
France (Lusignan)	>+360	>+360
Tanzania (Morogoro)	>+360	>+360
USA (Ames)	+269	>+360
Wheat	[CO <sub>2</sub> ]	[CO <sub>2</sub> ]
Argentina (Balcarce)	+117	+358
Australia (Wongan Hills)	0	+59
India (New Delhi)	+112	+278
Netherlands (Wageningen)	+83	+222
Rice	[CO <sub>2</sub> ]	[CO <sub>2</sub> ]
Philippines (Los Baños)	+131	>+360
China (Nanjing)	+30	+160
India (Ludhiana)	+348	+360
Japan (Shizukuishi)	+1	+87

concentration for a given crop in a given site. A probability of 0.5 indicates no evidence of yield loss or yield gain.

The higher the CO<sub>2</sub> concentration, the lower the risk of yield loss (Fig. 5). The yield loss probability curves show different patterns between crops, sites and temperature change. For maize (Fig. 5A and B), yield loss probabilities are almost always higher than 0.5 (with one exception in Ames (USA) for a temperature increase of +2 °C, and CO<sub>2</sub> concentrations higher than 250 ppm). For wheat (Fig. 5C and D), yield loss probabilities become lower than 0.5 in all sites for at least some of the considered CO<sub>2</sub> concentrations. In one site (Wongan Hills in Australia), these probabilities are even systematically lower than 0.5 for all tested concentration when the temperature increase is equal to +2 °C (Fig. 5C). These results show that, for wheat, a majority of crop models predict that an increase of CO<sub>2</sub> concentration can outweigh the negative effect of a temperature increase. Results obtained for rice are highly variable across sites (Fig. 5E and F). In some sites, the yield loss probability is lower than 0.5 for relatively low levels of CO<sub>2</sub> concentration increases, whereas the probability remains higher than 0.5 for all tested CO<sub>2</sub> concentrations in other sites (Los Baños in Philippines, and Ludhiana in India).

The curves presented in Fig. 5 were used to compute the thresholds of [CO<sub>2</sub>] increase required to obtain a probability of maize, wheat, and rice yield gain higher than 0.5 (i.e., more than 50% chance of yield gain). These thresholds were computed for two values of temperature increase (+2 and +4 °C) and four sites per crop (Table 1). As expected, these thresholds are all higher than +360 ppm for maize (the highest CO<sub>2</sub> concentration increase considered in this study) with one exception (Ames in USA with a temperature increase of +2 °C). For wheat, the thresholds range from +0 to +117 ppm and from +59 ppm to +358 ppm for a temperature increase of +2 °C and +4 °C respectively. The thresholds take intermediate values for rice (Table 1).

#### 4. Discussion and conclusions

Our study shows how yields simulated by ensembles of process-based dynamic crop models can be summarized by statistical models (meta-models) that are based on random coefficient regressions. These statistical models describe the between-crop model variability of the simulated yield data using probability distributions. They can be used to compute key simulated quantities such as mean yield loss, percentiles of yield loss, and probabilities of

yield loss as functions of temperature change and CO<sub>2</sub> concentration change. These statistical models are helpful for analyzing risk of yield loss due to climate change factors at the locations where the original simulations were conducted, but for a higher number of climate scenarios and without the need of re-running the original ensemble of process-based models.

Regression models with random parameters were previously used in a meta-analysis on the effect of climate change on crop yields (Wilcox and Makowski, 2014), where yield data were extracted from published papers, and random parameters were used to describe the between-site variability of the yield response to climate change factors. In these simulation experiments, the results varied not only due to different locations, crop management and climate data, but also due to the use and parameterization of different individual crop models (Wilcox and Makowski, 2014; Challinor et al., 2014). Contributing variability to model uncertainty and natural variability, an important distinction for using such results for decision making (Lehmann and Rillig, 2014), was not possible in the studies by Wilcox and Makowski (2014) and Challinor et al. (2014) due to the non-systematic variation in models, model parameters, sites, and climate scenarios considered. In our study, the use of standardized protocols for each model and applied across the three crops allowed a clear separation of causes of impact variability due to models, sites and climate factors. Therefore, in our study, the random parameter distributions applied do not describe the between-site variability but the variability among process-based crop models. The statistical models proposed in this study thus correspond to meta-models emulating ensembles of complex crop models.

An important advantage of our meta-models is that they handle the interpolation between temperature levels and between CO<sub>2</sub> concentration levels in a standardized manner. Our meta-models can thus be used to calculate temperature and [CO<sub>2</sub>] thresholds leading to yield loss or yield gain, and these thresholds can help agricultural and climate scientists to identify the climate change scenarios that are likely to lead to yield gains and yield losses. The capabilities of the meta-models were illustrated using three yield datasets generated by the AgMIP model intercomparison studies for maize, wheat, and rice (Asseng et al., 2013; Bassu et al., 2014; Li et al., 2015). Our results show that, for wheat, climate change has a 50% chance to result in a yield gain if [CO<sub>2</sub>] increases by at least +117 ppm (depending on the site) (i.e., [CO<sub>2</sub>] = 507 ppm) and if temperature concurrently increases by +2 °C (Table 1). Required levels of [CO<sub>2</sub>] increase were found to be much higher for maize, and intermediate for rice. It is important to mention that the [CO<sub>2</sub>] thresholds for yield gain/loss are related to the baseline scenario considered for simulating yield data. The use of other baselines (characterized by different temperature regimes) may lead to different thresholds. We do not advise using the meta-models for temperature and CO<sub>2</sub> concentration changes beyond the intervals considered in the original protocols. These intervals are already large (from 0 to +9 °C for temperature changes and from 0 to +360 ppm) and there is thus little practical interest in considering more extreme temperature and [CO<sub>2</sub>] increases. Moreover, there is no guarantee that the meta-models will perform correctly for more extreme climate scenarios.

Our meta-models can be used to quantify the effects of temperature and [CO<sub>2</sub>] on yields, their interactions, and their variability between sites and between all the considered crop models. They thus constitute powerful tools for exploring process-based crop model responses to climate factors. Results obtained here partly confirm those obtained by Wilcox and Makowski (2014) who found that the effects of high CO<sub>2</sub> concentrations outweighed the effects of increasing temperature (up to +2 °C), leading to increasing yields of wheat. However, the CO<sub>2</sub> concentration threshold leading to a yield gain is smaller in our study (from 390 to 507 ppm,

depending on the sites) than in Wilcox and Makowski (2014) (640 ppm in average). This difference is partly due to the fact that Wilcox and Makowski (2014) considered a higher number of sites located in many different countries whereas only four sites were considered here per crop.

Our meta-models also show that the divergence among the maize, wheat, and rice crop models (and therefore the uncertainty in the simulated results) increases as a function of temperature (the higher the temperature change, the higher the between-crop model variability) due to model uncertainties. The effect of CO<sub>2</sub> concentration on the variability among crop models is much smaller. This is consistent with the individual crop results reported by Asseng et al. (2013), Bassu et al. (2014), and Li et al. (2015). This however does not necessarily mean that models capture the effects of increased CO<sub>2</sub> better than the effects of increased temperature: rather it could indicate that models use similar approaches to simulate the effect of elevated CO<sub>2</sub>. Therefore, caution is required when models are used for impact assessments with late century climate scenarios where temperature changes are expected to be large. Multi-model applications (Martre et al., 2015) and model improvements with field experimental data (Asseng et al., 2015) are needed to reduce model uncertainties for such assessments.

One of the main interests of the proposed statistical model lies in its ability to describe the between-crop-model variability using probability distribution functions. The estimated parameter values of this statistical model are linked to the chosen ensemble of process-based crop models used to simulate the yield outputs. The use of a different set of crop models may change the fitted responses. For example, if one uses an ensemble of crop models all showing similar responses to temperature and to CO<sub>2</sub> concentration, the estimated variances of the random coefficients will be close to zero and the fitted yield probability distribution will be narrow and peaked. On the other hand, if the crop models included in the ensemble show contrasting responses, the estimated variances of the random coefficients will be high, and the fitted yield probability distribution will be less peaked and more flat.

The results presented in this paper are only valid for the locations and the range of climate conditions considered for fitting the statistical model. The simulated yield responses are thus site-specific. In future work, the meta-models presented here could be extended in two different ways. First, they could be applied to a dataset including simulations obtained for a larger number of sites. It may then be possible to improve the meta-models by including covariables describing site characteristics (e.g., soil type, agricultural practices) in order to explain the dominant causes of the between-site variability. Second, our meta-models could be extended in order to describe the between-year variability of yields for different climate scenarios. This could be achieved by using a more complex statistical model in order to describe yearly yield values and their distributions. Simulated yearly yields are likely to be correlated across models, and more sophisticated probability distributions thus need to be considered in order to provide a realistic description of the data.

## Acknowledgements

G.J. O'Leary was supported by the Victorian *Department of Economic Development Jobs, Transport and Resources*, the Australian *Department of Agriculture*. S. Bassu, P. Bertuzzi, G. De Sanctis, J.-L. Durand, D. Makowski, P. Martre, D. Ripoche and D. Wallach were partly supported by the INRA ACCAF meta-program. S. Gayler was supported by a grant from the Ministry of Science, Research and Arts of Baden-Württemberg (AZ Zu33-721.3-2) and the Helmholtz Centre for Environmental Research – UFZ, Leipzig. J. E. Olesen was funded through the FACCE MACSUR project by Innovation Fund

Denmark. F. Ewert and C. Angulo received support from the FACCE MACSUR project (031A103B) funded through the German Federal Ministry of Education and Research (2812ERA115). C. Biernath was funded through the Helmholtz project 'REKLIMRegional Climate Change: Causes and Effects' Topic 5: 'Chemistry-climate interactions on global to regional scales'. El Teixeira was supported by the Royal Society of New Zealand and the Climate Change Impacts and Implications for New Zealand (CCII) project. R. C. Izaurralde and J. R. Williams were funded by Texas AgriLife Research and Extension, Texas A&M University. C. O. Stöckle was funded through USDA National Institute for Food and Agriculture award 32011-680002-30191. C. Müller was funded through the FACCE MACSUR project by the German Ministry for Education and Research (BMBF, 031A103B). K.C. Kersebaum and C. Nendel were supported by FACE MACSUR project funded through the German Federal Office for Agriculture and Food (2812ERA147). C. Nendel received support by BMBF via the CARBIOCL research project (01LL0902M).

Y. Zhu and L. Tang were supported by the National High-Tech Research and Development Program of China (2013AA100404) and the Priority Academic Program Development of Jiangsu Higher Education Institutions in China (PAPD).

## References

- Akaike, H., 1973. *Information theory and an extension of the maximum likelihood principle*. In: Petrov, B.N., Csaki, F. (Eds.), *2nd International Symposium on Information Theory*. Akademia Kiado, Budapest, pp. 267–281.
- Asseng, S., Ewert, F., Rosenzweig, C., Jones, J.W., Hatfield, J.L., Ruane, A.C., Boote, K.J., Thorburn, P.J., Rotter, R.P., Cammarano, D., Brisson, N., Basso, B., Martre, P., Aggarwal, P.K., Angulo, C., Bertuzzi, P., Biernath, C., Challinor, A.J., Doltra, J., Gayler, S., Goldberg, R., Grant, R., Heng, L., Hooker, J., Hunt, L.A., Ingwersen, J., Izaurralde, R.C., Kersebaum, K.C., Müller, C., Naresh Kumar, S., Nendel, C., O'Leary, G., Olesen, J.E., Osborne, T.M., Palosuo, T., Priesack, E., Ripoche, D., Semenov, M.A., Shcherbak, I., Steduto, P., Stockle, C., Stratonovitch, P., Streck, T., Supit, I., Tao, F., Travasso, M., Waha, K., Wallach, D., White, J.W., Williams, J.R., Wolf, J., 2013. *Uncertainty in simulating wheat yields under climate change*. *Nat. Clim. Change* 3, 827–832.
- Asseng, S., Ewert, F., Martre, P., Rötter, R.P., Lobell, D.B., Cammarano, D., Kimball, B.A., Ottman, M.J., Wall, G.W., White, J.W., Reynolds, M.P., Alderman, P.D., Prasad, P.V.V., Aggarwal, P.K., Anothai, J., Basso, B., Biernath, C., Challinor, A.J., De Sanctis, G., Doltra, J., Fereres, E., Garcia-Vila, M., Gayler, S., Hoogenboom, G., Hunt, L.A., Izaurralde, R.C., Jabloun, M., Jones, C.D., Kersebaum, K.C., Koehler, A.K., Müller, C., Naresh Kumar, S., Nendel, C., O'Leary, G., Olesen, J.E., Palosuo, T., Priesack, E., Eysa Rezaei, E., Ruane, A.C., Semenov, M.A., Shcherbak, I., Stöckle, C., Stratonovitch, P., Streck, T., Supit, I., Tao, F., Thorburn, P., Waha, K., Wang, E., Wallach, D., Wolf, J., Zhao, Z., Zhu, Y., 2015. *Rising temperatures reduce global wheat production*. *Nat. Clim. Change* 5, 143–147. <http://dx.doi.org/10.1038/nclimate2470>.
- Bassu, S., Brisson, N., Durand, J.-L., Boote, K., Lizaso, J., Jones, J.W., Rosenzweig, C., Ruane, A.C., Adam, M., Baron, C., Basso, B., Biernath, C., Boogaard, H., Conijn, S., Corbeels, M., Deryng, D., De Sanctis, G., Gayler, S., Grassini, P., Hatfield, J., Hoek, S., Izaurralde, C., Jongschaap, R., Kemanian, A., Kersebaum, K.C., Kumar, N., Makowski, D., Müller, C., Nendel, C., Priesack, E., Pravia, M.V., Soo Hyung, K., Sau, F., Shcherbak, I., Tao, F., Teixeira, E., Timlin, D., Waha, K., 2014. *How do various maize crop models vary in their responses to climate change factors?* *Glob. Change Biol.* 20, 2301–2320. <http://dx.doi.org/10.1111/gcb.12520>.
- Challinor, A.J., Watson, J., Lobell, D.B., Howden, S.M., Smith, D.R., Chhetri, N., 2014. *A meta-analysis of crop yield under climate change and adaptation*. *Nat. Clim. Change* 4, 287–291.
- Davidian, M., Giltinan, D.M., 1995. *Nonlinear Mixed Effects Models for Repeated Measurement Data*. Chapman and Hall.
- Hatfield, J.L., Boote, K.J., Kimball, B.A., Izaurralde, R.C., Ort, D., Thomson, A., Wolfe, D.W., 2011. *Climate impacts on agriculture: implications for crop production*. *Agron. J.* 103, 351–370.
- Lehmann, J., Rillig, M., 2014. *Distinguishing variability from uncertainty*. *Nat. Clim. Change* 4, 153. <http://dx.doi.org/10.1038/nclimate2133>.
- Li, T., Hasegawa, T., Yin, X., Zhu, Y., Boote, K.J., Adam, M., Bregaglio, S., Buis, S., Confalonieri, R., Fumoto, T., Gaydon, D., Marcaida, M., Nakagawa, H., Oriol, P., Ruane, A.C., Ruget, F., Singh, B., Singh, U., Tang, L., Tao, F., Wilkens, P., Yoshida, H., Zhang, Z., Bouman, B., 2015. *Uncertainties in predicting rice yield by current crop models under a wide range of climatic conditions*. *Glob. Change Biol.* 21, 1328–1341. <http://dx.doi.org/10.1111/gcb.12758>.
- Lobell, D.B., Schlenker, W., Costa-Roberts, J., 2011. *Climate trends and global crop production since 1980*. *Science* 333, 616–620.
- Martre, P., Wallach, D., Asseng, S., Ewert, F., Jones, J.W., Rötter, R.P., Boote, K.J., Ruane, A.C., Thorburn, P.J., Cammarano, D., Hatfield, J.L., Rosenzweig, C., Aggarwal, P.K., Angulo, C., Basso, B., Bertuzzi, P., Biernath, C., Brisson, N., Challinor, A.J., Doltra, J., Gayler, S., Goldberg, R., Grant, R.F., Heng, L., Hooker, J., Hunt, L.A., Ingwersen, J.C., Izaurralde, R.C., Kersebaum, K.C., Müller, C., Kumar,

- S.N., Nendel, C., O'Leary, G.J., Olesen, J.E., Osborne, T.M., Palosuo, T., Priesack, E., Ripoche, D., Semenov, M.A., Shcherbak, I., Steduto, P., Stöckle, C.O., Stratonovitch, P., Streck, T., Supit, I., Tao, F., Travasso, M., Waha, K., White, J.W., Wolf, J., 2015. Multimodel ensembles of wheat growth: many models are better than one. *Glob. Change Biol.* 21, 911–925, <http://dx.doi.org/10.1111/gcb.12768>.
- Pinheiro, J.C., Bates, D.M., 2000. *Mixed-Effects Models in S and S-PLUS*. Springer, New York.
- Rosenzweig, C., Jones, J.W., Hatfield, J.L., Ruane, A.C., Boote, K.J., Thorburn, P., Antle, J.-M., Nelson, G.C., Porter, C., Janssen, S., Asseng, S., Basso, B., Ewert, F., Wallach, D., Baigorria, G., Winter, J.M., 2013. *The Agricultural Model Intercomparison and Improvement Project (AgMIP): Protocols and pilot studies*. *Agric. For. Meteorol.* 170, 166–182.
- Semenov, M.A., Wolf, J., Evans, L.G., Eckersten, H., Iglesias, A., 1996. *Comparison of wheat simulation models under climate change. 2. Application of climate change scenarios*. *Clim. Res.* 7, 271–281.
- Tubiello, F., Ewert, F., 2002. *Simulating the effects of elevated CO<sub>2</sub> on crops: approaches and applications for climate change*. *Eur. J. Agron.* 18, 57–74.
- White, J.W., Hoogenboom, G., Kimball, B.A., Wall, G.W., 2011. *Methodologies for simulating impacts of climate change on crop production*. *Field Crop Res.* 124, 357–368.
- Wilcox, J., Makowski, D., 2014. *A meta-analysis of the predicted effects of climate change on wheat yields using simulation studies*. *Field Crop Res.* 156, 180–190.